


PBW SEX PHEROMONE TRAPS DATA 2017-18 IN PUNJAB (AS IT STANDS ON 30.3.2018)




Increase in Peak from previous year of the corresponding period is due to the inversion of Cotton stick heaps .The Maximum no of Moth catches per night were found in Districts Vehari 1.24 per night and Lodhran 0.50 per night respectively.

COTTON GINNING WASTE FOUND & DISPOSED OFF DURING 4TH WEEK OF MARCH 2018 COMPARED WITH CORRESPONDING PERIOD 2017



Total Cotton Waste Found and Disposed Off in 2017 in Current Week was Nil While in 2018 the total cotton waste found and disposed off is 160 Kg.


COTTON GINNING WASTE DISPOSED OFF DURING 2017- 18



The highest Cotton ginning waste disposed off during 2018 was 63,618 Kg in 2nd week of January and minimum was 160 kg in 4th week of March.

The highest Cotton ginning waste disposed off during 2017 was 27,625 Kg in 2nd week of January and minimum was 0 in 4th week of March.


COTTON GINNING WASTE FOUND DURING 2017- 18



The highest Cotton ginning waste Found during 2018 was 73,923 Kg in 2nd week of January and minimum was 160 Kg in 4th week of March.


The highest Cotton ginning waste Found during 2017 was 27,625 Kg in 2nd week of January and minimum was 0 in 4th week of March.

DISTRICT WISE COTTON GINNING WASTE FOUND & DISPOSED OFF DURING 4TH WEEK OF MARCH 2018




Cotton Ginning waste found and disposed of for the current week is 160 Kg in Lodhran District only.

DISTRICT WISE PBW INSPECTION IN COTTON FACTORIES WASTE IN PUNJAB DURING 4TH WEEK OF MARCH 2018



PBW Larvae were inspected in Cotton Ginning Waste in Lodhran District was 2.00 Per Kg waste for the Current Week

DISTRICT WISE PBW INSPECTION IN COTTON STICK HEAPS IN PUNJAB DURING 4TH WEEK OF MARCH 2018



The maximum PBW Larvae were found in Khanewal while the minimum were found in Rajanpur District for the current week