

PEST SITUATION ON COTTON CROP IN PUNJAB DURING 2ND WEEK OF JULY, 2023

Whitefly: On overall Punjab basis 6.97% above ETL spots have been observed during current week as compared to 4.74% % in last week and 21.81% during the same corresponding period of last year. And 88.41% below ETL spots have been observed during current week as compared to 89.53% in last week and 77.07% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of whitefly

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	SARGODHA	50.0	27	JARANWALA	6.2
2	KASUR	50.0	28	TANDLIANWALA	6.2
3	MIAN CHANNU	45.9	29	FAISALABAD	5.6
4	CHICHA WATNI	40.5	30	KOT CHUTTA	5.4
5	VEHARI	35.4	31	JHANG	5.1
6	KOTMOMIN	33.3	32	KAROR L.E.	4.5
7	BURE WALA	27.5	33	LAYYAH	4.5
8	KABIR WALA	22.2	34	JAHANIAN	4.4
9	MAILSI	20.0	35	CHISHTIAN	4.0
10	CHUNIAN	20.0	36	MUZUFAR GARH	4.0
11	RAJAN PUR	17.2	37	A.P SIAL	3.8
12	KHANEWAL	17.1	38	HASILPUR	3.8
13	SAHIWAL	14.8	39	R.Y.KHAN	3.8
14	ARIF WALA	13.6	40	SHORKOT	3.4
15	KALLOR KOT	10.3	41	GOJRA	3.4
16	PIR MAHAL	9.4	42	MULTAN	2.8
17	RENALA KHURD	9.1	43	MIANWALI	2.6
18	CHAK JUMRAH	9.1	44	JALAL PUR P.W	2.5
19	DARYA KHAN	8.7	45	BAHAWAL NAGAR	2.5
20	KEHROR PACCA	7.4	46	FORT ABBAS	2.1
21	D.G.KHAN	7.2	47	KOT ADDU	1.9
22	T.T.SINGH	7.1	48	KHAN PUR	1.8
23	SUMUNDRI	7.1	49	LIAQUAT PUR	1.7
24	OKARA	6.9	50	ALI PUR	1.7
25	PAK PATTAN	6.7	51	CHOBARA	1.7
26	JAM PUR	6.5	52	BAHAWAL PUR	1.6

Jassid: On overall Punjab basis 17.23% above ETL spots have been observed as compared to 14.07% in last week and same week of last year 15.90% and 80.54% below ETL spots have been observed during current week as compared to 83.12% in last week and 81.93% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of Jassid

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	KOTMOMIN	66.7	33	LAYYAH	16.4
2	KEHROR PACCA	40.7	34	A.P.EAST	16.0
3	BAHAWAL PUR	38.1	35	GOJRA	15.3
4	MULTAN	36.1	36	SADIQ ABAD	15.1
5	DUYAPUR	35.7	37	KALLOR KOT	13.8
6	SHUJABAD	35.6	38	KOT ADDU	13.5
7	LODHRAN	33.8	39	MINCHAN ABAD	13.3
8	JALAL PUR P.W	32.5	40	BURE WALA	12.5
9	D.G.KHAN	29.0	41	MAILSI	12.5
10	CHOBARA	28.3	42	ALI PUR	11.9
11	KAROR L.E.	27.3	43	HAROON ABAD	11.8

12	JAHANIAN	26.7	44	SUMUNDRI	10.7
13	CHISHTIAN	26.0	45	BHAKKAR	10.7
14	MUZUFAR GARH	26.0	46	OKARA	10.3
15	RAJAN PUR	25.9	47	JHANG	10.2
16	SAHIWAL	25.0	48	BAHAWAL NAGAR	10.0
17	KHANEWAL	24.4	49	PIPLAN	10.0
18	KOT CHUTTA	24.3	50	T.T.SINGH	9.5
19	R.Y.KHAN	22.6	51	PIR MAHAL	9.4
20	YAZMAN	22.0	52	CHAK JUMRAH	9.1
21	VEHARI	20.8	53	DARYA KHAN	8.7
22	MANKERA	20.0	54	A.P SIAL	7.7
23	HASILPUR	18.9	55	SAHIWAL	7.4
24	FORT ABBAS	18.8	56	PAK PATTAN	6.7
25	TAUNSA	18.3	57	JARANWALA	6.2
26	ARIF WALA	18.2	58	TANDLIANWALA	6.2
27	RENALA KHURD	18.2	59	JATOI	5.4
28	MIAN CHANNU	18.0	60	SHORKOT	3.4
29	KHAN PUR	17.9	61	ROJHAN	3.0
30	KABIR WALA	17.8	62	FAISALABAD	2.8
31	JAM PUR	17.4	63	MIANWALI	2.6
32	LIAQUAT PUR	17.2			

Thrips: On overall Punjab basis 7.07% above ETL spots have been observed as compared to 4.35% in last week and same week of last year i.e. 3.86%. And 90.57% below ETL spots have been observed during current week as compared to 92.90% in last week and 92.49% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of Thrips

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	MINCHAN ABAD	10.0	26	T.T.SINGH	2.4
2	BAHAWAL NAGAR	10.0	27	KABIR WALA	2.2
3	LODHRAN	8.8	28	A.P.EAST	2.0
4	LIAQUAT PUR	8.6	29	A.P SIAL	1.9
5	ALI PUR	8.5	30	SAHIWAL	1.9
6	KAROR L.E.	7.6	31	JHANG	1.7
7	SADIQ ABAD	7.5	32	JARANWALA	1.5
8	BURE WALA	7.5	33	TANDLIANWALA	1.5
9	MAILSI	7.5	34	T.T.SINGH	2.4
10	KHANEWAL	7.3	35	KABIR WALA	2.2
11	KHAN PUR	7.1	36	A.P.EAST	2.0
12	JATOI	7.1	37	A.P SIAL	1.9
13	SHUJABAD	6.7	38	SAHIWAL	1.9
14	CHOBARA	6.7	39	JHANG	1.7
15	JAM PUR	6.5	40	JARANWALA	1.5
16	MIANWALI	6.5	41	TANDLIANWALA	1.5
17	MULTAN	5.6	42	T.T.SINGH	2.4
18	GOJRA	5.1	43	KABIR WALA	2.2
19	JALAL PUR P.W	5.0	44	A.P.EAST	2.0
20	MIAN CHANNU	4.9	45	A.P SIAL	1.9
21	JAHANIAN	4.4	46	SAHIWAL	1.9
22	DARYA KHAN	4.3	47	JHANG	1.7
23	HASILPUR	3.8	48	JARANWALA	1.5
24	PAK PATTAN	3.3	49	TANDLIANWALA	1.5
25	LAYYAH	3.0	50	T.T.SINGH	2.4

Mealy Bug: On overall Punjab basis 0.83% spots (all considered above ETL) of Mealy Bug have been observed as compared to 0.23% in last week and 3.61% spots during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of mealy bug

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	RAJAN PUR	12.1	8	DUYAPUR	2.4
2	ALI PUR	6.8	9	VEHARI	2.1
3	MANKERA	6.7	10	KHAN PUR	1.8
4	JALAL PUR P.W	5.0	11	JATOI	1.8
5	MUZUFAR GARH	4.0	12	MIAN CHANNU	1.6
6	MULTAN	2.8	13	BAHAWAL PUR	1.6
7	MAILSI	2.5	14	ROJHAN	1.5

Mites: On overall Punjab basis 0.17% above ETL spots has been observed during current week as compared to 0.00% in last week and 0.00% during the same corresponding period of last year. And 3.22% below ETL spots have been observed during current week as compared to 2.19% in last week and 0.88% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of mites

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	MAILSI	5.0	3	DUYAPUR	2.4
2	R.Y.KHAN	3.7736			

Dusky Cotton Bug: On overall Punjab basis 0.03% above ETL spots have been observed during current week as compared to 0.00% in last week and 0.08% during the same corresponding period of last year. And 15.07% below ETL spots have been observed during current week as compared to 13.22% in last week and 20.12% during the same corresponding period of last year.

Tehsil wise percentage of hot spots of dusky cotton bug

S. NO.	TEHSIL	%AGE
1	SHUJABAD	2.2

Armyworm: On overall Punjab basis 0.20% spots (all considered above ETL) of Armyworm have been observed as compared to 0.03% in last week and 0.24% spots during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of army worm

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	ALI PUR	3.4	4	JAHANIAN	2.2
2	MULTAN	2.8	5	FORT ABBAS	2.1
3	MAILSI	2.5			

Pink Boll Worm: On overall Punjab basis 0.30% above ETL spots have been observed as compared to 0.43% in last week and 0.08% same corresponding period of the last year. And 8.60% below ETL spots have been observed as compared to 8.34% in last week and 4.30% during same corresponding period of last year.

Tehsil wise percentage of hot spots of pink boll worm

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	JATOI	3.6	5	R.Y.KHAN	1.9
2	RAJAN PUR	3.4	6	ALI PUR	1.7
3	FORT ABBAS	2.1	7	BAHAWAL PUR	1.6
4	A.P.EAST	2.0			

American Boll Worm: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and 0.04% during same corresponding period of last year. And 0.53% below ETL spots have been observed as compared to 0.36% in last week and 0.08% during same corresponding period of last year.

Tehsil wise percentage of hot spots of American Boll Worm

No hot spot find yet.

Spotted Boll Worm: On overall Punjab basis 0.00% above ETL spots have been observed as compared to 0.00% in last week and 0.04% during same corresponding period of last year & 0.27% below ETL spots have been observed as compared to 0.10% in last week and 0.04% during same corresponding period of last year.

Tehsil wise percentage of hot spots of Spotted Boll Worm

No hot spot find yet.

CLCuV: On overall Punjab basis 0.27% spots (all considered above ETL) of CLCV have been observed as compared to 0.07% in last week and 19.88% during the same corresponding period of the last year.

Tehsil wise percentage of hot spots of CLCuV

S. NO.	TEHSIL	%AGE	S.NO.	TEHSIL	%AGE
1	BAHAWAL PUR	6.3	4	ALI PUR	1.7
2	VEHARI	2.1	5	LODHRAN	1.3
3	LIAQUAT PUR	1.7			

FORECAST FOR THE NEXT WEEK

Meteorological data of the current week 2022-2023

Districts	2023				2022			
	Temperature °C		R.H%	Rainfall (mm)	Temperature °C		RH%	Rainfall (mm)
	Max.	Min.			Max.	Min.		
Multan	39.4	30.3	60.8	0.0	38.7	29.9	61.6	0.0
Khanewal	40.3	28.4	56.7	0.0	39.4	31.0	60.7	0.0
Vehari	39.0	30.3	52.6	0.6	39.0	31.3	58.9	0.4
Lodhran	40.3	25.0	64.9	0.3	37.5	28.7	72.1	3.7
Sahiwal	35.5	26.2	46.7	27.6	38.4	28.2	76.1	44.0
Pakpattan	36.0	27.2	45.3	11.2	37.1	28.1	69.5	34.5
Okara	35.3	27.4	46.1	9.6	35.2	27.5	71.2	34.0
Bahawalpur	38.1	28.1	63.0	8.0	39.6	26.9	68.9	13.0
Bahawalnagar	37.4	28.2	61.9	7.0	41.7	28.5	41.7	27.0
R.Y.Khan	39.5	28.6	48.9	0.0	38.1	27.8	68.4	0.0
D.G. Khan	40.5	30.0	57.2	0.0	42.1	25.5	55.3	6.0
Muzaffar Garh	41.5	31.5	56.0	2.0	43.0	28.0	53.5	4.0
Rajanpur	40.3	30.3	57.5	3.0	37.3	28.3	58.2	4.0
Layyah	43.6	31.3	56.5	8.0	46.3	29.3	49.3	6.0
TOT/AVG	39.05	28.77	55.29	77.3	39.53	28.50	61.81	176.6

Weather forecast for next 7 day in cotton zone

Division	Dated	17/7	18/7	19/7	20/7	21/7	22/7	23/7
Multan	Max.Temp.	40	42	41	37	40	42	42
	Min.Temp.	30	33	31	32	30	32	34
	Humidity %	40	35	39	48	40	35	36
Sahiwal	Max.Temp.	28	40	40	36	39	40	39
	Min.Temp.	29	30	30	30	28	30	31
	Humidity %	46	41	43	50	47	41	44
Bahawalpur	Max.Temp.	40	41	40	37	37	41	40
	Min.Temp.	32	33	31	32	31	32	34
	Humidity %	40	36	38	50	52	39	39
D.G. Khan	Max.Temp.	40	43	42	40	39	42	42
	Min.Temp.	29	33	32	32	29	31	34
	Humidity %	41	32	33	41	47	38	36
Average Cotton Region	Max.Temp.	39.64						
	Min.Temp.	31.25						
	Humidity %	40.96						

Source: timeanddate.com

Summary of weather forecast

Overall weather for seven days in cotton zone during next week will remain hot, sunny & shiny.

Forecast of Sucking Pests:

Whitefly: This pest flourishes best in hot dry climate with optimum temperature 35-42 °C with relative humidity below 50%. Maximum lethal temperature for this pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. Keeping in view above all, it is predicted that population of this pest will increase during the coming week.

Jassid: This pest flourishes best in hot and humid climate with optimum temperature 35-40 °C with relative humidity above 70%. Maximum lethal temperature for this pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. Keeping in view above all, it is predicted that population of this pest will increase during the coming week.

Thrips: This pest flourishes best in hot dry climate with optimum temperature 35-40 °C with relative humidity below 60%. Maximum lethal temperature for this pest is 45-51 °C. . The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. In view above all, it is predicted that population of this pest will increase during the coming week.

Mealy Bug: This pest flourishes best in moderate climate with optimum temperature 35-40 °C with relative humidity above 60-80%. Maximum lethal temperature for this pest is 45-51 °C. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. Keeping in view above all, it is

predicted that population of this pest will sustain as such on cotton crop during the coming week.

Mites: This pest flourishes best in dry climate with optimum temperature 28-40 °C with relative humidity above 40-50%. Maximum lethal temperature for this pest is 45 °C. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. Keeping in view above all, it is predicted that population of this pest will sustain as such during the coming week.


Dusky Cotton Bug: This pest flourishes best in moderate climate with optimum temperature 28-40 °C with relative humidity above 50-80%. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. In view above all, it is predicted that population of this pest may sustain during the coming week.

Pink Bollworm: This pest flourishes best at moderate climate with optimum temperature 32-40 °C with relative humidity above 60-80%. Maximum lethal temperature for this pest is 45-53 °C. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. It is predicted that population of this pest will increase on cotton during the coming week.

Moth Catches in Pheromone Traps during 2nd Week of July, 2023

Moth Catches in Pheromone Traps During 2nd Week of July, 2023			
District	Traps Installed	Moth catches/Trap/week	Av. Moth catches/Trap/night
Multan	6	24	0.57
Khanewal	8	41	0.73
Vehari	6	26	0.62
Lodhran	6	40	0.95
Sahiwal	4	21	0.75
Pakpattan	4	23	0.82
Okara	4	21	0.75
Bahawalpur	8	50	0.89
Bahawalnagar	10	49	0.70
R.Y.Khan	8	47	0.84
D.G.Khan	6	8	0.19
Rajanpur	6	9	0.21
M.Garh	8	16	0.29
Layyah	6	14	0.33
Bhakkar	8	29	0.52
Mianwali	6	20	0.48
Faisalabad	6	46	1.10
TT. Singh	6	5	0.12
Jhang	4	17	0.61

Note. 3 moth catches on 3 consecutive nights in a trap is considered as ETL



American Bollworm: This pest flourishes best at dry & moderate climate with optimum temperature 32-40 °C with relative humidity above 50-70%. Maximum lethal temperature for this pest is 45-53 °C. The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. It is predicted that population of this pest will remain sustain on cotton crop during the coming week.

Moth catches on light traps during 2nd week of July, 2023

Moth Catches on Light Traps/Trap/night During 2nd Week of July, 2023				
District	Traps Installed	A.B.W	S.B.W	Armyworm
Multan	3	1.3	0.0	3.1
Khanewal	4	1.4	0.0	3.6
Vehari	3	0.6	0.0	3.7
Lodhran	3	1.6	0.0	4.1
Sahiwal	2	0.7	0.0	4.3
Pakpattan	2	0.4	0.0	3.7
Okara	3	0.7	0.0	3.1
Bahawalpur	4	1.4	0.7	2.1
Bahawalnagar	5	1.1	0.5	2.6
R.Y.Khan	4	0.6	0.7	2.6
D.G.Khan	3	0.9	0.3	3.4
M.Garh	4	0.0	0.0	3.0
Rajan Pur	3	0.6	0.6	2.9
Layyah	3	0.0	0.0	2.1
Note. 3 moth catches of a pest on 3 consecutive nights in a trap is considered as ETL				

Spotted Bollworm: This pest flourishes best at mild & humid climate with optimum temperature 30-40 °C with relative humidity above 60-100%. Maximum lethal temperature for this pest is 45-53 °C. . The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. Due to cultivation of more BT varieties, it is predicted that population of this pest will remain as such on cotton crop in the coming week.

Armyworm: This pest flourishes best in mild & humid climate with optimum temperature 30-40 °C with relative humidity below 70-100%. . The current weather conditions on overall Punjab basis (temp. Max. 39.05 °C, Min. 28.77 °C with R. humidity 55.29%) & forecast for the next week temp. Max. 39.64 °C, Min. 31.25°C with humidity 40.96. Keeping in view above all, it is predicted that population of this pest will sustain during the coming week.

CLCuV: The incidence of this disease is expected to sustain as such on cotton crop during the next week.

RECOMMENDATIONS

WHITEFLY MANAGEMENT

1. Conduct regular pest scouting of Whitefly.
2. Control the pest without delay when population reaches ETL i.e. nymphs or adults or both 5 per leaf.
3. Control weeds in and outside the field especially near water channels and fallow lands.

Alternate host plants of whitefly

Crops	Vegetables	Orchards	Weeds/ Ornamentals	Trees
Sunflower	Okra	Citrus	Gardenia	Shisham
Tobacco	Brinjal	Litchi	Mako	Shareen
	Cucurbits	Pomegranate	Maina	
	Tomato	Ber (<i>Zizyphus</i>)	Karund	
	Cabbage	Guava	Lehli	
	Cauliflower	Mulberry		
	Peas	Papaya		
	Potato			
	Onion			
	Spinach			

JASSID MANAGEMENT

1. Eradicate weeds.
2. Conduct Pest Scouting at least twice a week.

3. Use *Chrysoperla* Cards @ 80-90 cards per acre having 20-25 eggs on each card which are available at AARI Faisalabad, Sahiwal, Okara and Vehari.
(Repeat this process after every two weeks).

THRIPS MANAGEMENT

1. Conduct Pest Scouting regularly at least twice a week.
2. Avoid intercropping of cotton crop with vegetables.
3. Control weeds in and outside the field.

MEALY BUG MANAGEMENT

1. Visit the fields daily to detect the infestation for effective management of mealy bug well in time.
2. Keep field and water courses etc. free from weeds especially Itsit & Hazaardani, kangi booti, aksin etc.
3. To avoid its shifting from one place to another, keep the horticultural nurseries free from its infestation. Spray such infested nurseries before shifting the plants.
4. Avoid water stress to the plants.
5. Avoid repeated visits of workers from infested fields to healthy fields.
6. Apply the localized chemical spray on infected along with its 2 meter surrounding area.

Alternate host plants of cotton mealy bug

Crops	Vegetables	Ornamentals	Weeds	Orchards
Sunflower	Okra	China Rose	Hazar Dani	Citrus
Tobacco	Brinjal	Huddle	Amarantus	Mulberry
Jantar	Tomato	Cotton Rose	Bhakra	Ficus
Sesame	Chillies	Gulchain	Mako	Ber
	Pumpkin	Lantana	Sueda	
		Din Ka Raja	Itsit	
		Rat Ki Rani	Karund	

		Anthorium	Aksen	
		Gul-e-Daudi	Bathu	
		Gainda	Puth Kanda	
			Kanghi	

MANAGEMENT OF DUSKY COTTON BUG

1. Keep the special observation/ pest scouting when the boll opening starts.
2. Use of soft contact poison on the above ETL.

MANAGEMENT OF ARMY WORM

1. Keep the fields clean from weeds especially Its it which is a preferred host of armyworm.
2. Observe the field on regular basis and under close observation to assess its attack and treat the patch at earliest if necessary.
3. Avoid sowing of Jantar near cotton fields.
4. Regular pest scouting at least twice a week.
5. Hand picking and destruction of egg masses at initial stage as initially its attack starts in patches and is clearly visible.

CLCuV MANAGEMENT

1. Use CLCuV tolerant varieties for sowing.
2. Keep Cotton fields free from all kinds of weeds.
3. Eradicate other alternate host plants of CLCuV and dispose them off carefully.
4. Thinning out and destruction of Virus affected plants.
5. Remove CLCuV affected plants in less than 60 DAS (Days after sowing) crop.
6. Hoeing after each irrigation or rain at wattar condition till canopies permit.

7. Apply post emergence Glyphosate for both broad and narrow leaved weeds and Glyphosate for grassy weeds with shield and use Flat Fan Nozzle, if needed.
8. Irrigate the fields when needed keeping in view weather conditions e.g. temperature, rainfall, soil water holding capacity and plant need.
9. Balanced use of Fertilizer i.e. N.P.K. along with foliar growth promoters if necessary.
10. Keep transmission vector (whitefly) at the low ebb

Alternate host plants of CLCuV vector (whitefly)

Crops	Vegetables	Ornamentals	Weeds
Sunflower	Okra	Gurhal	Leh
Melon	Brinjal	Chambeli	Lehli
Tobacco	Chillies		Mako
	Tomato		Maina
	Potato		Karund/Bathu
	Cucumber		Gardenia
			Hazardani
			Rattanjot
			Sun Kukra

Economic threshold levels of cotton pests

Insect pests	Economic threshold levels
Jassid	1 Adult or Nymph per leaf
Whitefly	5 Adults or Nymphs or both per Leaf
Thrips	8-10 Adults or Nymphs per Leaf
Mealy bug	0 (on appearance)
Mites	On damage appearance
Spotted Boll worm	3 Larvae/ 25 plants.
Pink Boll worm	5 Larvae /100 bolls.
American boll worm	5 brown eggs or 3 small larvae or both 5 per 25 plants on non Bt varieties. 2 larvae of 2 nd instar per 25 plants on Bt varieties.
Army worm	0 (On appearance)
Aphid	Spray on visible damage on top terminals.
Dusky Cotton bug	10 per leaf/bud

